

Kollagen in Gefahr: Wenn die Architektur des Körpers den Halt verliert

Wie Spike mehr denn je und fortlaufend über Perizyten, Neutrophile Elastase Kollagen abbaut und durch Mikrothromben die Sauerstoffversorgung untergräbt

Ursula Ehrhorn

www.ursula-ehrhorn.com

Linktr.ee/doc.ehrhorn

Instagram | Youtube | E-Learning Academy

Literatur

- [1] Fernández-de-las-Peñas C et al. (2025): SARS-CoV-2 Spike Protein and Long COVID – Part 1. *Viruses* 17(5):617. <https://doi.org/10.3390/v17050617>
- [2] Fernández-de-las-Peñas C et al. (2025): SARS-CoV-2 Spike Protein and Long COVID – Part 2. *Viruses* 17(5):619. <https://doi.org/10.3390/v17050619>
- [3] Hoffmann M et al. (2020): SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2. *Cell* 181(2):271–280. <https://doi.org/10.1016/j.cell.2020.02.052>
- [4] Huang L et al. (2024): Persistence of spike protein at the skull-meninges-brain axis. *Cell Host & Microbe* 36(10):1433–1447. <https://doi.org/10.1016/j.chom.2024.09.013>
- [5] Ackermann M et al. (2020): Pulmonary Vascular Endothelialitis, Thrombosis, and Angiogenesis in Covid-19. *NEJM* 383:120–128. <https://doi.org/10.1056/NEJMoa2015432>
- [6] Sigrist CJ et al. (2020): A potential role for integrin cell-adhesion molecules in SARS-CoV-2 infection. *Antiviral Research* 177:104759. <https://doi.org/10.1016/j.antiviral.2020.104759>
- [7] Bhatt DL et al. (2022): Receptor binding domain of SARS-CoV-2 is a functional α -integrin agonist. *J Mol Biol* 434(9):167371. PMID: 35441172.
- [8] Yurchenco PD (2011): Basement Membranes: Cell Scaffoldings and Signaling Platforms. *Cold Spring Harbor Perspect Biol* 3:a004911. <https://doi.org/10.1101/cshperspect.a004911>
- [9] Tandon R et al. (2025): Heparan sulfate proteoglycans remodel SARS-CoV-2 spike conformation. *Front Cell Infect Microbiol* 15:1552116. <https://doi.org/10.3389/fcimb.2025.1552116>
- [10] Papayannopoulos V (2018): Neutrophil extracellular traps in immunity and disease. *Nat Rev Immunol* 18:134–147. <https://doi.org/10.1038/nri.2017.105>
- [11] Veras FP et al. (2020): SARS-CoV-2-triggered neutrophil extracellular traps mediate COVID-19 pathology. *J Exp Med* 217(12):e20201129. <https://doi.org/10.1084/jem.20201129>
- [12] Middleton EA et al. (2020): Neutrophil extracellular traps contribute to immunothrombosis in COVID-19 ARDS. *Blood* 136(10):1169–1179. <https://doi.org/10.1182/blood.2020007008>
- [13] Nicolai L et al. (2022): NETosis and Neutrophil Extracellular Traps in COVID-19. *Front Immunol* 13:838011. <https://doi.org/10.3389/fimmu.2022.838011>
- [14] Xiao Y et al. (2024): Neutrophil extracellular traps mediate neurological dysfunction in COVID-19. *Neural Regen Res* 19(8). <https://doi.org/10.4103/1673-5374.390964>
- [15] Hazeldine J, Lord JM (2023): Neutrophil extracellular traps and long COVID. *Front Immunol* 14:1254310. <https://doi.org/10.3389/fimmu.2023.1254310>
- [16] Nyström S, Hammarström P (2022): Amyloidogenesis of SARS-CoV-2 Spike Protein. *J Am Chem Soc* 144(20):8945–8950. <https://doi.org/10.1021/jacs.2c03925>
- [17] Pretorius E et al. (2021): SARS-CoV-2 spike protein S1 induces fibrin(ogen) resistant to fibrinolysis. *Biosci Rep* 41(8):BSR20210611. <https://doi.org/10.1042/BSR20210611>
- [18] Westman H, Hammarström P, Nyström S (2025): SARS-CoV-2 Spike Protein Amyloid Fibrils Impair Fibrin Formation and Fibrinolysis. *Biochemistry* 64(24):4818–4829. <https://doi.org/10.1021/acs.biochem.5c00550>
- [19] Bhatt DL et al. (2024): The Possible Mechanistic Basis of Individual Susceptibility to Spike Protein Injury. *J Thromb Haemost*. PMID:12213048.
- [20] Brumback BD et al. (2022): Preferential uptake of SARS-CoV-2 by pericytes potentiates vascular dysfunction. *Cardiovasc Res*. PMID: 35709328.
- [21] Brumback BD et al. (2022): Human Cardiac Pericytes Are Susceptible to SARS-CoV-2 Infection. *JACC Basic Transl Sci*. PMID:9473702.
- [22] Li P, Fan H (2023): Pericyte Loss in Diseases. *Cells* 12(15):1931. <https://doi.org/10.3390/cells12151931>
- [23] Sardu C et al. (2022): SARS-CoV-2 and vascular dysfunction: a growing role for pericytes. *Cardiovasc Res*. PMID:9494379.
- [24] He L et al. (2020): Pericyte-specific vascular expression of SARS-CoV-2 receptor ACE2. *bioRxiv*.

- [25] Varga Z et al. (2021): Rapid endothelitis and vascular damage characterize SARS-CoV-2 infection. PMID: 33908688.
- [26] Vuong JT et al. (2023): Post COVID-19 Large Vessel Vasculopathy. *Semin Thromb Hemost* 49(3):282–290. <https://doi.org/10.1055/s-0042-1758122>
- [27] Stikvoort B et al. (2023): Extracellular Matrix Remodeling in Vascular Disease. *Arterioscler Thromb Vasc Biol* 43(8):1406–1419.
- [28] Gu Z et al. (2020): Rapid, Nitric Oxide Synthesis-Dependent Activation of MMP-9 in Neutrophils. *Front Physiol* 11:9. <https://doi.org/10.3389/fphys.2020.00009>
- [29] Haljasmägi L et al. (2024): Extracellular matrix remodelling pathway in peripheral blood in COVID-19. *Front Immunol* 15:1379356. <https://doi.org/10.3389/fimmu.2024.1379356>
- [30] Avolio E et al. (2021): Emerging Role of Pericytes and Their Secretome in the Heart. *Cells* 10(3):548. <https://doi.org/10.3390/cells10030548>
- [31] Jin Y et al. (2022): Endothelial dysfunction in COVID-19: an overview of evidence, biomarkers, mechanisms and potential therapies. *Acta Pharmacol Sin* 43(5):1083–1096. <https://doi.org/10.1038/s41401-021-00753-7>
- [32] Luszczyk B et al. (2022): Role of the Extracellular Matrix in SARS-CoV-2 Infection. *Int J Mol Sci* 23(3):1152. <https://doi.org/10.3390/ijms23031152>
- [33] Teuwen LA et al. (2020): COVID-19: the vasculature unleashed. *Nat Rev Immunol* 20(7):389–391. <https://doi.org/10.1038/s41577-020-0343-0>
- [34] COVID-19 and cellular senescence. *Nat Rev Immunol*. 2022. <https://doi.org/10.1038/s41577-022-00785-2>
- [35] Chen S et al. (2023): Baicalin ameliorates endothelial dysfunction and platelet activation via inhibition of the Furin/TGFβ1/Smad3/TSP-1 pathway. *Eur J Pharmacol* 955:175715. <https://doi.org/10.1016/j.ejphar.2023.175715>
- [36] Sekheri M et al. (2022): The "Elastic Perspective" of SARS-CoV-2 Infection and the Role of Neutrophil Elastase in Vascular Damage. *Int J Mol Sci* 23(3):1576. PMID: 8835950.
- [36] Stark K et al. (2022): Pericyte-derived chemokines amplify neutrophil recruitment. *Nat Commun* 13:4422. PMID: 9371542.
- [37] Mustafa Z et al. (2021): Neutrophil Elastase and Proteinase 3 Cleavage Sites Are Adjacent to the Polybasic Sequence within the Proteolytic Sensitive Activation Loop of the SARS-CoV-2 Spike Protein. *ACS Omega* 6(10):7181–7185. <https://doi.org/10.1021/acsomega.1c00363>
- [38] Baig AM, Rosko S, Jaeger B, Gerlach J, Rausch H (2024): Unraveling the enigma of long COVID: novel aspects in pathogenesis, diagnosis, and treatment protocols. *Inflammopharmacology* 32(4):2075–2090. <https://doi.org/10.1007/s10787-024-01483-2>
- [39] Gerlach J, Baig AM, Fabrowski M, Viduto V (2023): The immune paradox of SARS-CoV-2: Lymphocytopenia and autoimmunity evoking features in COVID-19 and possible treatment modalities. *Rev Med Virol* 33(2):e2423. <https://doi.org/10.1002/rmv.2423>
- [40] Baig AM, Rosko S, Jaeger B, Gerlach J (2025): Strategic Inhibition of CHRM Autoantibodies: Molecular Insights and Therapeutic Potentials in Long COVID. *J Med Chem* 68(3):2089–2094. <https://doi.org/10.1021/acs.jmedchem.4c00691>
- [41] Kim et al. (2025): Prolonged immune activation in post-acute sequelae of SARS-CoV-2: neutrophil dynamics and therapeutic insights. *Exp Mol Med*. <https://doi.org/10.1038/s12276-025-01539-5>
- [42] Kell DB, Laubscher GJ, Pretorius E (2022): A central role for amyloid fibrin microclots in long COVID/PASC: origins and therapeutic implications. *Biochem J* 479(4):537–559. <https://doi.org/10.1042/BCJ20220016>
- [43] Pretorius E et al. (2022): Proteomics of fibrin amyloid microclots in long COVID/PASC shows entrapped pro-inflammatory molecules. *Cardiovasc Diabetol* 21:190. <https://doi.org/10.1186/s12933-022-01623-4>
- [44] Pretorius E et al. (2022): Prevalence of fibrin amyloid microclots and platelet pathology in Long COVID/PASC. *Cardiovasc Diabetol* 21:148. <https://doi.org/10.1186/s12933-022-01579-5>