

Jenseits von Abnehmspritze und reiner Kalorienrestriktion: Naturstoffe zur Modulation von GLP-1, Insulin und Sättigung

Kira Sidnal, M.Sc.

Eisenstraße 1

57482 Wenden | Deutschland

kira.sidnal@tisso.de

www.tisso.de

Literatur

- [1] Boix-Castejon M, Herranz-Lopez M, Perez Gago A, Olivares-Vicente M, Caturla N, Roche E, et al. Hibiscus and lemon verbena polyphenols modulate appetite-related biomarkers in overweight subjects: a randomized controlled trial. *Food Funct.* 2018;9(6):3173–84.
- [2] Capomolla AS, Janda E, Paone S, Parafati M, Sawicki T, Mollace R, et al. Atherogenic Index Reduction and Weight Loss in Metabolic Syndrome Patients Treated with A Novel Pectin-Enriched Formulation of Bergamot Polyphenols. *Nutrients.* 2019;11(6).
- [3] Schellenberg R, Zimmermann C, Drewe J, Hoexter G, Zahner C. Dose-dependent efficacy of the Vitex agnus castus extract Ze 440 in patients suffering from premenstrual syndrome. *Phytomedicine.* 2012;19(14):1325–31.
- [4] Thondre PS, Lightowler H, Ahlstrom L, Gallagher A. Mulberry leaf extract improves glycaemic response and insulaemic response to sucrose in healthy subjects: results of a randomized, double blind, placebo-controlled study. *Nutr Metab (Lond).* 2021;18(1):41.
- [5] Thomas JP, A.; Sivasadan, S. D.; Sreevallabhan, S.; Madhavamenon, K. I.; Mohanan, R. Clove bud (*Syzygium aromaticum* L.) polyphenol helps to mitigate metabolic syndrome by establishing intracellular redox homeostasis and glucose metabolism: A randomized, double-blinded, active-controlled comparative study. *Journal of Functional Foods.* 2022;98.
- [6] Noubiap JJ, Nansseu JR, Lontchi-Yimagou E, Nkeck JR, Nyaga UF, Ngouo AT, et al. Geographic distribution of metabolic syndrome and its components in the general adult population: A meta-analysis of global data from 28 million individuals. *Diabetes Res Clin Pract.* 2022;188:109924.
- [7] Schutte S, Eberhard S, Burger B, Hemmerling M, Rossol S, Stahmeyer JT. [Prevalence of metabolic syndrome : Analysis based on routine statutory health insurance data]. *Inn Med (Heidelb).* 2023;64(5):482–9.
- [8] Diabetes mellitus: Prävalenz (ab 18 Jahre): Robert Koch-Institut; 2025 (Available from: https://www.gbe.rki.de/DE/Themen/Gesundheitszustand/KoerperlicheErkrankungen/DiabetesMellitus/DiabetesPraevalenz18Plus/diabetesPraevalenz_node.html?darstellung=0&kennzahl=1&zeit=2024&geslecht=0&standardisierung=0).
- [9] Schienkiewitz AK, R.; Blume, M.; Mensink, G. B. M. Übergewicht und Adipositas bei Erwachsenen in Deutschland - Ergebnisse der Studie GEDA 2019/2020-EHIS: Robert Koch-Institut; 2022 (Available from: <https://edoc.rki.de/handle/176904/10231>).
- [10] Arnold SE, Arvanitakis Z, Macauley-Rambach SL, Koenig AM, Wang HY, Ahima RS, et al. Brain insulin resistance in type 2 diabetes and Alzheimer disease: concepts and conundrums. *Nat Rev Neurol.* 2018;14(3):168–81.
- [11] Samuel VT, Shulman GI. Mechanisms for insulin resistance: common threads and missing links. *Cell.* 2012;148(5):852–71.
- [12] Giovannucci E. Insulin, insulin-like growth factors and colon cancer: a review of the evidence. *J Nutr.* 2001;131(11 Suppl):3109S–20S.
- [13] Barr EL, Boyko EJ, Zimmet PZ, Wolfe R, Tonkin AM, Shaw JE. Continuous relationships between non-diabetic hyperglycaemia and both cardiovascular disease and all-cause mortality: the Australian Diabetes, Obesity, and Lifestyle (AusDiab) study. *Diabetologia.* 2009;52(3):415–24.
- [14] Jacobs E, Hoyer A, Brinks R, Kuss O, Rathmann W. Burden of Mortality Attributable to Diagnosed Diabetes: A Nationwide Analysis Based on Claims Data From 65 Million People in Germany. *Diabetes Care.* 2017;40(12):1703–9.
- [15] Bjornholt JV, Erikssen G, Aaser E, Sandvik L, Nitter-Hauge S, Jervell J, et al. Fasting blood glucose: an underestimated risk factor for cardiovascular death. Results from a 22-year follow-up of healthy nondiabetic men. *Diabetes Care.* 1999;22(1):45–9.
- [16] Meigs JB, Nathan DM, D'Agostino RB, Sr., Wilson PW, Framingham Offspring S. Fasting and postchallenge glycemia and cardiovascular disease risk: the Framingham Offspring Study. *Diabetes Care.* 2002;25(10):1845–50.
- [17] Li M, Chi X, Wang Y, Setrerrahmane S, Xie W, Xu H. Trends in insulin resistance: insights into mechanisms and therapeutic strategy. *Signal Transduct Target Ther.* 2022;7(1):216.
- [18] Lu X, Xie Q, Pan X, Zhang R, Zhang X, Peng G, et al. Type 2 diabetes mellitus in adults: pathogenesis, prevention and therapy. *Signal Transduct Target Ther.* 2024;9(1):262.

- [19] Pagliai G, Dinu M, Madarena MP, Bonaccio M, Iacoviello L, Sofi F. Consumption of ultra-processed foods and health status: a systematic review and meta-analysis. *Br J Nutr*. 2021;125(3):308–18.
- [20] Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, et al. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab*. 2019;30(1):67–77 e3.
- [21] Hamano S, Sawada M, Aihara M, Sakurai Y, Sekine R, Usami S, et al. Ultra-processed foods cause weight gain and increased energy intake associated with reduced chewing frequency: A randomized, open-label, crossover study. *Diabetes Obes Metab*. 2024;26(11):5431–43.
- [22] Preston JM, Iversen J, Hufnagel A, Hjort L, Taylor J, Sanchez C, et al. Effect of ultra-processed food consumption on male reproductive and metabolic health. *Cell Metab*. 2025;37(10):1950–60 e2.
- [23] Galdino-Silva MB, Almeida KMM, Oliveira ADS, Santos J, Macena ML, Silva DR, et al. A Meal with Ultra-Processed Foods Leads to a Faster Rate of Intake and to a Lesser Decrease in the Capacity to Eat When Compared to a Similar, Matched Meal Without Ultra-Processed Foods. *Nutrients*. 2024;16(24).
- [24] Touvier M. Health effects of ultra-processed food: uncovering causal mechanisms. *Nat Rev Endocrinol*. 2026;22(2):66–7.
- [25] Mao Z, Zhang W. Role of mTOR in Glucose and Lipid Metabolism. *Int J Mol Sci*. 2018;19(7).
- [26] Stephens BR, Granados K, Zderic TW, Hamilton MT, Braun B. Effects of 1 day of inactivity on insulin action in healthy men and women: interaction with energy intake. *Metabolism*. 2011;60(7):941–9.
- [27] Dirks ML, Wall BT, van de Valk B, Holloway TM, Holloway GP, Chabowski A, et al. One Week of Bed Rest Leads to Substantial Muscle Atrophy and Induces Whole-Body Insulin Resistance in the Absence of Skeletal Muscle Lipid Accumulation. *Diabetes*. 2016;65(10):2862–75.
- [28] Sondrup N, Termannsen AD, Eriksen JN, Hjorth MF, Faerch K, Klingenberg L, et al. Effects of sleep manipulation on markers of insulin sensitivity: A systematic review and meta-analysis of randomized controlled trials. *Sleep Med Rev*. 2022;62:101594.
- [29] Hackett RA, Steptoe A. Type 2 diabetes mellitus and psychological stress - a modifiable risk factor. *Nat Rev Endocrinol*. 2017;13(9):547–60.
- [30] Gröber. Mikronährstoffe: MWW Medizinisch Wissenschaftliche Verlagsgesellschaft; 2019.
- [31] Via M. The malnutrition of obesity: micronutrient deficiencies that promote diabetes. *ISRN Endocrinol*. 2012;2012:103472.
- [32] Lepretti M, Martucciello S, Burgos Aceves MA, Putti R, Lionetti L. Omega-3 Fatty Acids and Insulin Resistance: Focus on the Regulation of Mitochondria and Endoplasmic Reticulum Stress. *Nutrients*. 2018;10(3).
- [33] Wu J, Atkins A, Downes M, Wei Z. Vitamin D in Diabetes: Uncovering the Sunshine Hormone's Role in Glucose Metabolism and Beyond. *Nutrients*. 2023;15(8).
- [34] Kostov K. Effects of Magnesium Deficiency on Mechanisms of Insulin Resistance in Type 2 Diabetes: Focusing on the Processes of Insulin Secretion and Signaling. *Int J Mol Sci*. 2019;20(6).
- [35] Garofalo V, Condorelli RA, Cannarella R, Aversa A, Calogero AE, La Vignera S. Relationship between Iron Deficiency and Thyroid Function: A Systematic Review and Meta-Analysis. *Nutrients*. 2023;15(22).
- [36] Cruz KJC, de Oliveira ARS, Morais JBS, Severo JS, Mendes PMV, de Sousa Melo SR, et al. Zinc and Insulin Resistance: Biochemical and Molecular Aspects. *Biol Trace Elem Res*. 2018;186(2):407–12.
- [37] Corbin KD, Zeisel SH. Choline metabolism provides novel insights into nonalcoholic fatty liver disease and its progression. *Curr Opin Gastroenterol*. 2012;28(2):159–65.
- [38] Mishra S, Jain S, Agadzi B, Yadav H. A Cascade of Microbiota-Leaky Gut-Inflammation- Is it a Key Player in Metabolic Disorders? *Curr Obes Rep*. 2025;14(1):32.
- [39] Pan H, Guo J, Su Z. Advances in understanding the interrelations between leptin resistance and obesity. *Physiol Behav*. 2014;130:157–69.
- [40] Holst JJ. The physiology of glucagon-like peptide 1. *Physiol Rev*. 2007;87(4):1409–39.
- [41] Hardie DG. AMPK: a key regulator of energy balance in the single cell and the whole organism. *Int J Obes (Lond)*. 2008;32 Suppl 4:S7–12.
- [42] Zheng Z, Zong Y, Ma Y, Tian Y, Pang Y, Zhang C, et al. Glucagon-like peptide-1 receptor: mechanisms and advances in therapy. *Signal Transduct Target Ther*. 2024;9(1):234.
- [43] Rubino D, Abrahamsson N, Davies M, Hesse D, Greenway FL, Jensen C, et al. Effect of Continued Weekly Subcutaneous Semaglutide vs Placebo on Weight Loss Maintenance in Adults With Overweight or Obesity: The STEP 4 Randomized Clinical Trial. *JAMA*. 2021;325(14):1414–25.
- [44] He L, Wang J, Ping F, Yang N, Huang J, Li Y, et al. Association of Glucagon-Like Peptide-1 Receptor Agonist Use With Risk of Gallbladder and Biliary Diseases: A Systematic Review and Meta-analysis of Randomized Clinical Trials. *JAMA Intern Med*. 2022;182(5):513–9.
- [45] Filippatos TD, Panagiotopoulou TV, Elisaf MS. Adverse Effects of GLP-1 Receptor Agonists. *Rev Diabet Stud*. 2014;11(3–4):202–30.
- [46] Wilding JPH, Batterham RL, Davies M, Van Gaal LF, Kandler K, Konakli K, et al. Weight regain and cardiometabolic effects after withdrawal of semaglutide: The STEP 1 trial extension. *Diabetes Obes Metab*. 2022;24(8):1553–64.
- [47] Jastreboff AM, Aronne LJ, Ahmad NN, Wharton S, Connery L, Alves B, et al. Tirzepatide Once Weekly for the Treatment of Obesity. *N Engl J Med*. 2022;387(3):205–16.
- [48] Houmard JA, Tanner CJ, Yu C, Cunningham PG, Pories WJ, MacDonald KG, et al. Effect of weight loss on insulin sensitivity and intramuscular long-chain fatty acyl-CoAs in morbidly obese subjects. *Diabetes*. 2002;51(10):2959–63.

- [49] Clamp LD, Hume DJ, Lambert EV, Kroff J. Enhanced insulin sensitivity in successful, long-term weight loss maintainers compared with matched controls with no weight loss history. *Nutr Diabetes*. 2017;7(6):e282.
- [50] Trussardi Fayh AP, Lopes AL, Fernandes PR, Reischak-Oliveira A, Friedman R. Impact of weight loss with or without exercise on abdominal fat and insulin resistance in obese individuals: a randomised clinical trial. *Br J Nutr*. 2013;110(3):486–92.
- [51] Mohan R, Jose S, Mulakkal J, Karpinsky-Semper D, Swick AG, Krishnakumar IM. Water-soluble polyphenol-rich clove extract lowers pre- and post-prandial blood glucose levels in healthy and prediabetic volunteers: an open label pilot study. *BMC Complement Altern Med*. 2019;19(1):99.
- [52] Thondre PS, Butler I, Tammam J, Achebe I, Young E, Lane M, et al. Understanding the Impact of Different Doses of Reducose((R)) Mulberry Leaf Extract on Blood Glucose and Insulin Responses after Eating a Complex Meal: Results from a Double-Blind, Randomised, Crossover Trial. *Nutrients*. 2024;16(11).
- [53] Cui W, Luo K, Xiao Q, Sun Z, Wang Y, Cui C, et al. Effect of mulberry leaf or mulberry leaf extract on glycemic traits: a systematic review and meta-analysis. *Food Funct*. 2023;14(3):1277–89.
- [54] Chen S, Xi M, Gao F, Li M, Dong T, Geng Z, et al. Evaluation of mulberry leaves' hypoglycemic properties and hypoglycemic mechanisms. *Front Pharmacol*. 2023;14:1045309.
- [55] Lown M, Fuller R, Lightowler H, Fraser A, Gallagher A, Stuart B, et al. Mulberry-extract improves glucose tolerance and decreases insulin concentrations in normoglycaemic adults: Results of a randomised double-blind placebo-controlled study. *PLoS One*. 2017;12(2):e0172239.
- [56] Ding F, Wang Q, Xie C, Wang M, Zhang L, Gao M, et al. The impact of mulberry leaf extract at three different levels on reducing the glycemic index of white bread. *PLoS One*. 2023;18(8):e0288911.
- [57] Foster-Schubert KE, Overduin J, Prudom CE, Liu J, Callahan HS, Gaylinn BD, et al. Acyl and total ghrelin are suppressed strongly by ingested proteins, weakly by lipids, and biphasically by carbohydrates. *J Clin Endocrinol Metab*. 2008;93(5):1971–9.
- [58] Mammen RR, Natinga Mulakal J, Mohanan R, Maliakel B, Illathu Madhavamenon K. Clove Bud Polyphenols Alleviate Alterations in Inflammation and Oxidative Stress Markers Associated with Binge Drinking: A Randomized Double-Blinded Placebo-Controlled Crossover Study. *J Med Food*. 2018;21(11):1188–96.
- [59] Namazi N, Khodamoradi K, Khamechi SP, Heshmati J, Ayati MH, Larijani B. The impact of cinnamon on anthropometric indices and glycemic status in patients with type 2 diabetes: A systematic review and meta-analysis of clinical trials. *Complement Ther Med*. 2019;43:92–101.
- [60] Zarezadeh M, Musazadeh V, Foroumandi E, Keramati M, Ostadrahimi A, Mekary RA. The effect of cinnamon supplementation on glycemic control in patients with type 2 diabetes or with polycystic ovary syndrome: an umbrella meta-analysis on interventional meta-analyses. *Diabetol Metab Syndr*. 2023;15(1):127.
- [61] de Moura SL, Gomes BGR, Guillarducci MJ, Coelho OGL, Guimaraes NS, Gomes JMG. Effects of cinnamon supplementation on metabolic biomarkers in individuals with type 2 diabetes: a systematic review and meta-analysis. *Nutr Rev*. 2025;83(2):249–79.
- [62] Bakhach H, Nuffer M, Tall Bull S, Nuffer W. A Systematic Review Evaluating Cinnamon's Effects on Glucose Utilizing a Ranking System to Assess Bias and Study Quality. *J Med Food*. 2024;27(9):814–23.
- [63] Hohe tägliche Aufnahmemengen von Zimt: Gesundheitsrisiko kann nicht ausgeschlossen werden Bundesinstitut für Risikobewertung; 2006 [Available from: <https://www.bfr.bund.de/stellungnahme/hohe-taegliche-aufnahmemengen-von-zimt-gesundheitsrisiko-kann-nicht-ausgeschlossen-werden/>].
- [64] Fragen und Antworten zu Cumarin in Zimt und anderen Lebensmitteln: Bundesinstitut für Risikobewertung; 2012 [Available from: <https://www.bfr.bund.de/fragen-und-antworten/thema/fragen-und-antworten-zu-cumarin-in-zimt-und-anderen-lebensmitteln/>].
- [65] Zhao F, Pan D, Wang N, Xia H, Zhang H, Wang S, et al. Effect of Chromium Supplementation on Blood Glucose and Lipid Levels in Patients with Type 2 Diabetes Mellitus: a Systematic Review and Meta-analysis. *Biol Trace Elem Res*. 2022;200(2):516–25.
- [66] Burgerstein. *Handbuch Nährstoffe* 2023.
- [67] Gliozzi MC, C.; Musolino, V.; Palma, E.; Muscoli, C.; Vitale CG, S.; Muscianisi, G.; Janda, E.; Muscoli, S.; Romeo FR, S.; Mollace, R.; Walker, R.; Ehrlich, J.; Mollace V. The Effect of Bergamot-Derived Polyphenolic Fraction on LDL Small Dense Particles and Non Alcoholic Fatty Liver Disease in Patients with Metabolic Syndrome. *Advances in Biological Chemistry*. 2014;4.
- [68] Mollace V, Sacco I, Janda E, Malara C, Ventrice D, Colica C, et al. Hypolipemic and hypoglycaemic activity of bergamot polyphenols: from animal models to human studies. *Fitoterapia*. 2011;82(3):309–16.
- [69] Martínez-Rodríguez AM-O, M.; Vicente-Martínez, M.; Asencio-Mas, N.; Navarro, P.; Caturla, N.; Jones, J. . Effectiveness of a Polyphenol-Enriched Blend on Weight Management and Metabolic Syndrome-Related Parameters in Healthy Overweight Adults. *Appl Sci* 2024;14(9).
- [70] Herranz-Lopez M, Olivares-Vicente M, Boix-Castejon M, Caturla N, Roche E, Micol V. Differential effects of a combination of Hibiscus sabdariffa and Lippia citriodora polyphenols in overweight/obese subjects: A randomized controlled trial. *Sci Rep*. 2019;9(1):2999.

- [71] Marhuenda J, Perez S, Victoria-Montesinos D, Abellan MS, Caturla N, Jones J, et al. A Randomized, Double-Blind, Placebo Controlled Trial to Determine the Effectiveness a Polyphenolic Extract (*Hibiscus sabdariffa* and *Lippia citriodora*) in the Reduction of Body Fat Mass in Healthy Subjects. *Foods*. 2020;9(1).
- [72] Canfora EE, Jocken JW, Blaak EE. Short-chain fatty acids in control of body weight and insulin sensitivity. *Nat Rev Endocrinol*. 2015;11(10):577–91.
- [73] Li H, Zhang L, Li J, Wu Q, Qian L, He J, et al. Resistant starch intake facilitates weight loss in humans by reshaping the gut microbiota. *Nat Metab*. 2024;6(3):578–97.
- [74] Li S, Guerin-Deremaux L, Pochat M, Wils D, Reifer C, Miller LE. NUTRIOSE dietary fiber supplementation improves insulin resistance and determinants of metabolic syndrome in overweight men: a double-blind, randomized, placebo-controlled study. *Appl Physiol Nutr Metab*. 2010;35(6):773–82.
- [75] Jarrar AH, Stojanovska L, Apostolopoulos V, Feehan J, Bataineh MF, Ismail LC, et al. The Effect of Gum Arabic (*Acacia Senegal*) on Cardiovascular Risk Factors and Gastrointestinal Symptoms in Adults at Risk of Metabolic Syndrome: A Randomized Clinical Trial. *Nutrients*. 2021;13(1).
- [76] Guerin-Deremaux L, Pochat M, Reifer C, Wils D, Cho S, Miller LE. The soluble fiber NUTRIOSE induces a dose-dependent beneficial impact on satiety over time in humans. *Nutr Res*. 2011;31(9):665–72.
- [77] Mukai J, Tsuge Y, Yamada M, Otori K, Atsuda K. Effects of resistant dextrin for weight loss in overweight adults: a systematic review with a meta-analysis of randomized controlled trials. *J Pharm Health Care Sci*. 2017;3:15.
- [78] Parisi G, Bottona E, Carrara M, Cardin F, Faedo A, Goldin D, et al. Treatment effects of partially hydrolyzed guar gum on symptoms and quality of life of patients with irritable bowel syndrome. A multicenter randomized open trial. *Dig Dis Sci*. 2005;50(6):1107–12.