

Leistung entsteht in der Erholung

Funktionelle Diagnostik der mitochondrialen Adaptations- und Regenerationsfähigkeit

Prof. Dr. rer. nat. habil. Brigitte König
Breiter Weg 10A
39118 Magdeburg | Deutschland
mmd@mmd-labor.de
www.mmd-labor.de

Literatur

- [1] Smith JAB, Murach KA, Dyar KA, Zierath JR. Exercise metabolism and adaptation in skeletal muscle. *Nat Rev Mol Cell Biol.* 2023;24:607–632. PMID: 37225892.
- [2] Memme JM, Erlich AT, Phukan G, Hood DA. Exercise and mitochondrial health. *J Physiol.* 2021;599:803–817. PMID: 31674658.
- [3] Egan B, Zierath JR. Exercise metabolism and the molecular regulation of skeletal muscle adaptation. *Cell Metab.* 2013;17:162–184. PMID: 23395166.
- [4] Chacko BK et al. The Bioenergetic Health Index: a new concept in mitochondrial translational research. *Clin Sci (Lond).* 2014;127:367–373. PMID: PMC4202728.
- [5] Chacko BK et al. The Bioenergetic Health Index is a sensitive measure of oxidative stress in human monocytes. *Redox Biol.* 2016;8:43–50. PMID: 26748041.
- [6] Braganza A et al. Blood-based bioenergetics: an emerging translational and clinical tool. *Mol Aspects Med.* 2019;71:100835. PMID: PMC7031032.
- [7] Janssen JJE et al. Extracellular flux analyses reveal differences in mitochondrial PBMC metabolism between high-fit and low-fit females. *Am J Physiol Endocrinol Metab.* 2022. PMID: PMC8897018.
- [8] Gebhardt K et al. PBMCs mitochondrial respiration and its relation to immunity and inflammation in healthy older adults. 2025. PMID: PMC12475958.
- [9] Hedges CP et al. Peripheral blood mononuclear cells do not reflect skeletal muscle mitochondrial function or adaptation to high-intensity interval training in healthy young men. *J Appl Physiol.* 2019;126:454–461. PMID: 30571281.
- [10] Goodpaster BH, Sparks LM. Metabolic flexibility in health and disease. *Cell Metab.* 2017;25:1027–1036. PMID: 28467922.
- [11] San-Millán I, Brooks GA. Assessment of metabolic flexibility by means of measuring blood lactate, fat, and carbohydrate oxidation responses to exercise. *Sports Med.* 2018;48:467–479. PMID: 28623613.
- [12] Neuberger EWI et al. Physical activity specifically evokes release of cell-free DNA from granulocytes thereby affecting liquid biopsy. *Clin Epigenetics.* 2022;14:29. PMID: PMC8864902.
- [13] Haller N et al. Circulating, cell-free DNA as a marker for exercise load in intermittent sports. *PLoS One.* 2018;13:e0191915. PMID: PMC5784997.
- [14] Juškevičiūtė E et al. Cell-free DNA kinetics in response to muscle-damaging exercise in healthy individuals. *Eur J Appl Physiol.* 2024. PMID: 38875105.
- [15] Ohlsson L et al. Increased level of circulating cell-free mitochondrial DNA due to a single bout of strenuous physical exercise. *Eur J Appl Physiol.* 2020;120:897–905. PMID: 32088743.
- [16] Cerqueira É et al. Inflammatory effects of high and moderate intensity exercise: a systematic review. *Front Physiol.* 2020;10:1550. PMID: 31992987.
- [17] Nieman DC et al. Impact of a competitive marathon race on systemic cytokine and neutrophil responses. *Med Sci Sports Exerc.* 2003;35:66–75. PMID: 12569227.
- [18] Fonseca TR et al. Aerobic training modulates the increase in plasma concentrations of cytokines in response to a session of exercise. *J Immunol Res.* 2021;2021:1304139. PMID: 33510799.
- [19] Kleinert M et al. Exercise increases circulating GDF15 in humans. *Mol Metab.* 2018;9:187–191. PMID: 29398617.
- [20] Conte M et al. Altered GDF15 and FGF21 levels in response to strenuous exercise: a study in marathon runners. *Front Physiol.* 2020;11:550102. PMID: 33329017.